

MAGNETIC FIELDS IN MASSIVE STARS

S. Hubrig,¹

RESUMEN

Favor de proporcionar un resumen en español. If you cannot provide a spanish abstract, the editors will do this. Although indirect evidence for the presence of magnetic fields in high-mass stars is regularly reported in the literature, the detection of these fields remains an extremely challenging observational problem. We review the recent discoveries of magnetic fields in different types of massive stars and briefly discuss strategies for spectropolarimetric observations to be carried out in the future.

ABSTRACT

Although indirect evidence for the presence of magnetic fields in high-mass stars is regularly reported in the literature, the detection of these fields remains an extremely challenging observational problem. We review the recent discoveries of magnetic fields in different types of massive stars and briefly discuss strategies for spectropolarimetric observations to be carried out in the future.

Key Words: **STARS: CIRCUMSTELLAR MATTER — STARS: MAGNETIC FIELDS — STARS: OSCILLATION — STARS: POLARISATION**

1. INTRODUCTION

An increasing number of observations of hot stars provide indirect evidence that magnetic fields must be present in those stars: Cyclic behaviour on a rotational timescale observed in the UV wind lines (e.g., Henrichs et al. 2005), the presence of narrow X-ray emission lines (Cohen et al. 2003), non-thermal radio emission (Bieging et al. 1989), etc. In spite of numerous indirect evidence only very few direct magnetic field detections have been reported so far. Magnetic fields are accessible through the Zeeman effect: The Zeeman components of spectral lines are polarised and thus permit magnetic fields to be measured even in rapidly rotating massive stars where rotation broadening, etc., prevents the resolution of Zeeman components.

2. RECENT SEARCHES

Currently, direct measurements are achieved only in two O stars, θ^1 Ori C and HD 191612 with B_{eff} values of a few hundred Gauss (Donati et al. 2002, 2006). To study the incidence of magnetic fields in O stars, we recently obtained high S/N spectropolarimetric observations of eleven O stars with FORS1 at the VLT with a typical accuracy of the field determination of about 30–70 G. However, no evidence of a magnetic field was found, leading to the conclusion that large scale, dipole like magnetic fields are not widespread among O-type stars.

Among early B-type stars, a magnetic field has been discovered in the B0.2V star τ Sco and in one of the hottest β Cephei stars, the B0.7IV star ξ^1 CMa. Two more β Cephei stars, β Cep (Henrichs et al. 2000) and V2053 Oph (Neiner et al. 2003a), with spectral types B2III and B2IV, exhibit a weak magnetic field. The star ξ^1 CMa has the largest magnetic field of up to 300 G (Hubrig et al. 2006), whereas the magnetic field for the other three stars is much weaker with corresponding B_{eff} values less than 100 G. A magnetic field of the order of a few hundred Gauss has recently been discovered for slowly pulsating B-type stars (Neiner et al. 2003b, Hubrig et al. 2006). The effect of the magnetic field on the oscillation properties of β Cephei stars and slowly pulsating B-type stars has not been studied in detail yet. In Figs. 1 and 2 we present the most recent magnetic field observations of the hottest β Cephei star ξ^1 CMa.

Another type of massive stars, rapidly rotating Be stars, lose mass and initially accumulate it in a rotating circumstellar disk. Much of the mass loss is in the form of outbursts and so additional mechanisms such as the beating of nonradial pulsation modes or magnetic flares must be at work. Previously, only two Be stars have been found as weekly magnetic with B_{eff} values less than 100 G (Neiner et al. 2003c, Hubrig et al. 2006). Our recent search for magnetic fields in 15 Be stars with FORS1 at the VLT revealed that out of our sample five Be stars present circular polarisation signatures in lines

¹European Southern Observatory, Casilla 19001, Santiago 19, Chile (shubrig@eso.org).




Fig. 1. Stokes V/I spectra of the β Cephei star ξ^1 CMa around the H β line. The thickness of the plotted line corresponds to the uncertainty of the polarisation measurements determined from photon noise.




Fig. 2. Stokes V/I spectra of the β Cephei star ξ^1 CMa around the He II line at 4121 Å.

formed in the circumstellar environment (Hubrig et al. 2007). Furthermore, for a few stars we measured a field of the order of 100 G, changing its field polarity every 8–9 minutes. Clearly, time series are needed to study these local transient magnetic fields. As it has already been discussed by Maheswaran (2003) in the framework of his Magnetic Rotator Wind-

Disk Model, magnetic fields can consist of flux loops that emerge from lower latitudes and thread the disk around the Be star.

3. DISCUSSION

Magnetic fields are indeed present in hot massive stars. Potential progress in their study may come from achieving better accuracy of the measurements of magnetic fields or from the detailed studies of polarised line profiles, in the case that the magnetic fields of massive stars have structures significantly more complex than those of classical Ap and Bp stars.

The Hanle effect is a relatively new magnetic diagnostic in stellar astrophysics, but can probably be used to study circumstellar magnetic fields. This effect produces a depolarisation of the scattered radiation and a rotation of the plane of polarisation (e.g., Ignace et al. 2004). A program for such observations is currently allocated at the VLT with FORS 1 and we are expecting to receive the data in the middle of 2007.

REFERENCES

- Bieging, J. H., Abbott, D. C., & Churchwell, E. B. 1989, *ApJ*, 340, 518
- Cohen, D. H., et al. 2003, *ApJ*, 586, 495
- Donati, J.-F., Babel, J., Harries, T. J., Howarth, I. D., Petit, P., & Semel, M. 2002, *MNRAS*, 333, 55
- Donati, J.-F., Howarth, I. D., Bouret, J.-C., Petit, P., Catala, C., & Landstreet, J. 2006, *MNRAS*, 365, 6
- Henrichs, H. F., et al. 2000, in *ASP Conference Series*, Vol. 214, “The Be Phenomenon in Early-Type Stars”, eds. Smith, M. A., & Henrichs, H. F., p. 324
- Henrichs, H. F., Schnerr, R. S., & Ten Kulve, E. 2005, in *ASP Conference Series*, Vol. 337, “The Nature and Evolution of Disks Around Hot Stars”, eds. Ignace, R. & Gayley, K. G., p. 114
- Hubrig, S., Briquet, M., Schöller, M., De Cat, P., Mathys, G., & Aerts, C. 2006, *MNRAS*, 369, 61
- Hubrig, S., et al. 2007, *in preparation*
- Ignace, R., Nordsieck, K. H., & Cassinelli, J. P. 2004, *ApJ*, 609, 1018
- Maheswaran, M. 2003, *ApJ*, 592, 1156
- Neiner, C., et al. 2003a, *A&A*, 411, 565
- Neiner, C., et al. 2003b, *A&A*, 406, 1019
- Neiner, C., et al. 2003c, *A&A*, 409, 275